MAP 7

Globigerina bulloides

Absolute abundance

Summer	Fall	Winter	Spring	(no. of specimens per 1000m³)
○	□	△	✡	Absent
●	■	▲	✱	1-99
●	■	▲	✱	100-999
●	■	▲	✱	1000-9999
●	■	▲	✱	10,000 and greater

MAP 8

Globigerina bulloides

Relative abundance

Summer	Fall	Winter	Spring	(percent of total population)
○	□	△	✡	Absent
●	■	▲	✱	0.1-4.9
●	■	▲	✱	5.0-9.9
●	■	▲	✱	10.0-19.9
●	■	▲	✱	20.0 and greater

Globigerina bulloides
×42



MAP 9

Globorotalia truncatulinoides

Absolute abundance

Summer	Fall	Winter	Spring	(no. of specimens per 1000m³)
○	□	△	✡	Absent
●	■	▲	✱	1-99
●	■	▲	✱	100-999
●	■	▲	✱	1000-9999
●	■	▲	✱	10,000 and greater

MAP 10

Globorotalia truncatulinoides

Relative abundance

Summer	Fall	Winter	Spring	(percent of total population)
○	□	△	✡	Absent
●	■	▲	✱	0.1-4.9
●	■	▲	✱	5.0-9.9
●	■	▲	✱	10.0-19.9
●	■	▲	✱	20.0 and greater

Globorotalia truncatulinoides
×62



MAP 11

Globorotalia inflata

Absolute abundance

Summer	Fall	Winter	Spring	(no. of specimens per 1000m³)
○	□	△	✡	Absent
●	■	▲	✱	1-99
●	■	▲	✱	100-999
●	■	▲	✱	1000-9999
●	■	▲	✱	10,000 and greater

MAP 12

Globorotalia inflata

Relative abundance

Summer	Fall	Winter	Spring	(percent of total population)
○	□	△	✡	Absent
●	■	▲	✱	0.1-4.9
●	■	▲	✱	5.0-9.9
●	■	▲	✱	10.0-19.9
●	■	▲	✱	20.0 and greater

Globorotalia inflata
×68



MAP 13

Globigerinita glutinata

Absolute abundance

Summer	Fall	Winter	Spring	(no. of specimens per 1000m³)
○	□	△	✡	Absent
●	■	▲	✱	1-99
●	■	▲	✱	100-999
●	■	▲	✱	1000-9999
●	■	▲	✱	10,000 and greater

MAP 14

Globigerinita glutinata

Relative abundance

Summer	Fall	Winter	Spring	(percent of total population)
○	□	△	✡	Absent
●	■	▲	✱	0.1-4.9
●	■	▲	✱	5.0-9.9
●	■	▲	✱	10.0-19.9
●	■	▲	✱	20.0 and greater

Globigerinita glutinata
×71

Austral seasons are indicated

COMPILED BY A. W. H. BE

AMERICAN GEOGRAPHICAL SOCIETY—1969